



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let O be the given point, AB the given side. Let $OA=a$, $OB=b$, $OC=c$, $OD=d$, $AB=e$, $FA=x$, $AD=z$, $DG=y$, $\angle OFA=\theta$.

$$\text{Then } a^2 = (y+z)^2 + x^2 - 2x(y+z)\cos\theta \dots (1),$$

$$b^2 = (y+z)^2 + (x+e)^2 - 2(x+e)(y+z)\cos\theta \dots (2),$$

$$c^2 = y^2 + (x+e)^2 - 2y(x+e)\cos\theta \dots (3),$$

$$d^2 = y^2 + x^2 - 2xy\cos\theta \dots (4).$$

From (1) and (2),

$$y+z = x\cos\theta \pm \sqrt{a^2 - x^2\sin^2\theta} = (x+e)\cos\theta$$

$$\pm \sqrt{b^2 - (x+e)^2\sin^2\theta} \dots (5).$$

From (3) and (4),

$$y = (x+e)\cos\theta \pm \sqrt{c^2 - (x+e)^2\sin^2\theta} = x\cos\theta \pm \sqrt{d^2 - x^2\sin^2\theta} \dots (6).$$

$$\text{Let } 4b^2e^2 = A^2, (a^2 - e^2 - b^2) = B, 2ce = C, d^2 - e^2 - c^2 = D.$$

Then (5) and (6) become after reduction

$$A^2 - B^2 = [4Be(e+x) + A^2 + 4e^2(e+x)^2]\sin^2\theta \dots (7),$$

$$C^2 - D^2 = [4De(e+x) + C^2 + 4e^2(e+x)^2]\sin^2\theta \dots (8).$$

Eliminating $\sin^2\theta$ we get

$$4e^2(A^2 - B^2 - C^2 + D^2)(e+x)^2 + 4e[D(A^2 - B^2) - B(C^2 - D^2)](e+x) = B^2C^2 - A^2D^2 \dots (9).$$

Equation (9) determines x . Then (7) or (8) determines θ , (3) or (4) determines y , (1) or (2) determines z , and the parallelogram is determined in all respects.

228. Proposed by O.E. GLENN, A.M., Fellow in Mathematics, University of Pennsylvania, Philadelphia, Pa.


Given a point O without a circle S ; two arbitrary lines through O cut S in the points A, A' , and B, B' , respectively. Prove, by pure geometry, that the four circles through $OAR, OBR, OA'R', OB'R'$, respectively, intersect in points collinear with O ; R and R' being points upon S arbitrarily chosen.

No solution has been received.

CALCULUS.

179. Proposed by B. F. FINKEL, A. M., Drury College, Springfield, Mo.

Discuss the integrals of the equation $x(1-x)w'' + [1 - (a+b+1)x]w' - abw = 0$ in the vicinities $x=0$, and $x=1$; indicating the form for the latter vicinity when $a+b=1$. Also when $1-a-b$ is an integer k . [From Forsyth's *Linear Differential Equations*, Ex. 6, p. 103].



Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

This is the differential equation of the the well known hypergeometric series

$$w=1+\frac{ab}{1^2}x+\frac{a(a+1)b(b+1)}{1^2.2^2}x^2+\frac{a(a+1)(a+2)b(b+1)(b+2)}{1^2.2^2.3^2}x^3+\dots$$

discussed in Chapter VI of Forsyth's *Differential Equations*.

If $x=0$, $w=1$; if $x>1$, w is divergent; if $x<1$, w is convergent; if $x=1$, $1-a-b=k$, a positive integer, w is convergent; if $x=1$, $1-a-b=k$, a negative integer or zero, w is divergent; if $x=1$, $a+b=1$ and w is divergent.

MECHANICS.

167. Proposed by EDWIN S. CRAWLEY, Ph.D., Professor of Mathematics in the University of Pennsylvania

An anchor ring or torus is produced by the revolution of a circle of radius r , the center of the revolving circle describing a circle of radius R . A quadrant of the torus is cut by two planes through the center of the ring perpendicular to each other and perpendicular to the plane of revolution. Determine the limiting value of the ratio $R:r$, so that when the quadrant thus formed is placed with one of its ends in coincidence with a horizontal plane it will rest in that position.

IV. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let the circle radius r revolve around the line GH at a distance R from its center, through an angle $2\theta=\frac{1}{2}\pi$. Let $OM=x$, $MP=y$, $MN=dx$.

The figure $PQpq$ will, by revolution, generate a quadrantal-cylindrical shell whose volume is ultimately $2y\pi x dx$. The center of gravity of this shell will be on the axis of X at a distance $\frac{xs\sin\theta}{\theta}=\frac{2x\sqrt{2}}{\pi}$ from O .

$$\begin{aligned}\therefore \bar{x} &= \frac{\int_{R-r}^{R+r} \frac{2x\sqrt{2}}{\pi} \cdot 2y\pi x dx}{\int_{R-r}^{R+r} 2y\pi x dx} = \frac{2\sqrt{2}}{\pi} \cdot \frac{\int_{R-r}^{R+r} x^2 y dx}{\int_{R-r}^{R+r} xy dx} \\ &= \frac{2\sqrt{2}}{\pi} \cdot \frac{\int_{R-r}^{R+r} x^2 \sqrt{[r^2 - (x-R)^2]} dx}{\int_{R-r}^{R+r} x \sqrt{[r^2 - (x-R)^2]} dx} = \frac{(4R^2 + r^2)\sqrt{2}}{2\pi R}.\end{aligned}$$

$$\therefore FE = \frac{(4R^2 + r^2)\sqrt{2}}{3\pi R} \text{ or } FD = \frac{4R^2 + r^2}{2\pi R}.$$

Also $FK=R$, $DK=r$. By the conditions of the problem,

$$R-r = \frac{4R^2 + r^2}{2\pi R}. \quad \therefore \frac{R}{r} = \frac{\pi + \sqrt{(\pi^2 + 2\pi - 4)}}{2\pi - 4} = 2.9.$$

